

REMARK ON THE CALABI FLOW WITH BOUNDED CURVATURE

BY GÁBOR SZÉKELYHIDI

Abstract. In this short note we prove that if the curvature tensor is uniformly bounded along the Calabi flow and the Mabuchi energy is proper, then the flow converges to a constant scalar curvature metric.

Let (M, ω) be a compact Kähler manifold. For any Kähler potential φ , such that $\omega_\varphi = \omega + \sqrt{-1}\partial\bar{\partial}\varphi$ is positive, Aubin's I -functional [1] is defined by

$$I(\varphi) = \int_M \varphi(\omega^n - \omega_\varphi^n).$$

Our main result is the following.

THEOREM 1. *Given constants $K > 0$ and $\alpha \in (0, 1)$, there is a $C > 0$ with the following property. If $\omega_\varphi = \omega + \sqrt{-1}\partial\bar{\partial}\varphi$ satisfies $|Rm(\omega_\varphi)| < K$ and $I(\varphi) < K$, then $\omega_\varphi > C^{-1}\omega$ and $\|\omega_\varphi\|_{C^{1,\alpha}(\omega)} < C$.*

This result should be compared with Chen–He [4, Theorem 5.1], where only a bound on the Ricci curvature is assumed, but instead of $I(\varphi)$ the C^0 -norm of φ is assumed to be bounded. Note that in contrast with the result in [4], our proof is by contradiction and it does not give explicit control of the constant C . It would be interesting to obtain bounds on C depending on K and the geometry of (M, ω) . In Example 7 we will show that the assumption of a bound on $I(\varphi)$ is sharp in a certain sense.

The proof of Theorem 1 relies on two ingredients. One is the ϵ -regularity statement for harmonic maps, as was used in Ruan [17], and the other is some properties of plurisubharmonic functions and their Lelong numbers, taken from Guedj–Zeriahi [10]. We will review these in Section 1.

Our main application of the theorem is to the Calabi flow. This is the fourth order parabolic flow

$$\frac{\partial}{\partial t}\varphi = S(\omega_\varphi) - \hat{S},$$

introduced by Calabi [2], where $S(\omega_\varphi)$ is the scalar curvature of ω_φ and \hat{S} is its average. In Chen–He [4] it was shown that the flow exists as long as the Ricci curvature of the metrics remains bounded. In general little is known about the behavior of the Calabi flow, but there are many result in special cases, e.g. [5, 6, 9, 20]. In this paper we study the flow under the simplifying assumption that the curvature remains uniformly bounded for all time. The Kähler–Ricci flow has been studied previously under the same assumption (see e.g. [16, 21, 23]), the goal being to relate convergence of the flow to some algebro-geometric stability condition. The Calabi flow poses extra difficulties, since the diameter is not apriori bounded and collapsing can occur, as can be seen in the examples in [20]. Recently Huang [11, 12] has studied the flow on toric manifolds, and our result can be seen as extending some of his work to general Kähler manifolds. The following is a direct consequence of Theorem 1.

THEOREM 2. *Suppose that the Mabuchi energy is proper on the class $[\omega]$, and the curvature remains uniformly bounded along the Calabi flow with initial metric ω . The flow then converges exponentially fast to a constant scalar curvature metric in the Kähler class $[\omega]$.*

We will review the notion of properness of the Mabuchi energy in Section 1. The same proof can be used to prove a similar result under the assumption that the “modified” Mabuchi energy is proper, with the limit being an extremal metric, but we will not discuss this.

Acknowledgements. I would like to thank Valentino Tosatti and Hongnian Huang for useful comments.

1. Background material.

Plurisubharmonic functions and Lelong numbers. First we summarize the relevant ideas from [17]. The basic observation is that the identity map $\iota : (M, \omega) \rightarrow (M, \omega_\varphi)$ is harmonic. The energy density of ι is given by

$$e(\varphi) = \Lambda_\omega(\omega_\varphi) = \Delta\varphi + n.$$

The ϵ -regularity estimate of Schoen–Uhlenbeck [18] for harmonic maps states (Proposition 2.1 in [17]):

PROPOSITION 3. *Suppose that $|Rm(\omega_\varphi)| < K$. There exists an $\epsilon > 0$ depending on ω and K , such that if $r > 0$ and $x \in M$ satisfy*

$$(1) \quad r^{2-2n} \int_{B_r(x)} e(\varphi) \omega^n < \epsilon,$$

then

$$\sup_{B_{r/2}(x)} e(\varphi) < \frac{4r^{-2n}}{\epsilon} \int_{B_r(x)} e(\varphi) \omega^n < 4r^{-2}.$$

The key observation in [17] is that the expression in (1) also appears in the definition of the Lelong number of a plurisubharmonic function at x . We write

$$PSH(M, \omega) = \{\varphi \in L^1(M) : \varphi \text{ is upper semicontinuous,} \\ \text{and } \omega + \sqrt{-1}\partial\bar{\partial}\varphi \geq 0\}.$$

If $\varphi \in PSH(M, \omega)$ and $x \in M$, then the Lelong number [13] $\nu(\varphi, x)$ of φ at x is defined to be

$$\nu(\varphi, x) = \lim_{r \rightarrow 0} c_n r^{2-2n} \int_{B_r(x)} \sqrt{-1}\partial\bar{\partial}\varphi \wedge \omega^{n-1},$$

where c_n is a normalizing constant.

We now review the relevant results in [10]. Recall that

$$\mathcal{E}(M, \omega) \subset PSH(M, \omega)$$

is defined to be the set of $\varphi \in PSH(M, \omega)$, such that

$$\lim_{j \rightarrow \infty} (\omega + \sqrt{-1}\partial\bar{\partial}\varphi_j)^n (\varphi \leq -j) = 0,$$

where $\varphi_j = \max\{\varphi, -j\}$. This is a natural class of plurisubharmonic functions, on which the complex Monge–Ampère operator is well-defined. For us their most important property is Corollary 1.8 from [10]:

PROPOSITION 4. *Any $\varphi \in \mathcal{E}(M, \omega)$ has zero Lelong number at every $x \in M$.*

An important subset of $\mathcal{E}(M, \omega)$ consists of the elements of finite energy, $\mathcal{E}^1(M, \omega)$, defined by

$$\mathcal{E}^1(M, \omega) = \{\varphi \in \mathcal{E}(M, \omega) : \varphi \in L^1(M, \omega_\varphi^n)\}.$$

For elements in $\mathcal{E}^1(M, \omega)$ let us write

$$E(\varphi) = - \int_M \varphi \omega_\varphi^n.$$

Corollary 2.7 in [10] states:

PROPOSITION 5. *Suppose that $\varphi_j \in \mathcal{E}^1(M, \omega)$ is a sequence converging to φ in $L^1(M)$ such that $\varphi_j \leq 0$ and $E(\varphi_j)$ is uniformly bounded. Then $\varphi \in \mathcal{E}^1(M, \omega)$.*

The Mabuchi functional. The Mabuchi functional [14] is a functional

$$\mathcal{M} : [\omega] \rightarrow \mathbf{R}$$

on the Kähler class $[\omega]$, which is most easily defined by its variation. If $\omega_t = \omega + \sqrt{-1}\partial\bar{\partial}\varphi_t$, then

$$\frac{d}{dt}\mathcal{M}(\omega_t) = \int_M \varphi_t(\hat{S} - S(\omega_t))\omega_t^n,$$

where \hat{S} is the average scalar curvature. One can normalize so that $\mathcal{M}(\omega) = 0$ for a fixed reference metric $\omega \in [\omega]$. It is clear from the variation that constant scalar curvature metrics are the critical points of \mathcal{M} . For us the most important notion is that of properness.

DEFINITION 6. The Mabuchi energy is *proper* on the class $[\omega]$, if there is an increasing function $f : \mathbf{R} \rightarrow \mathbf{R}$ with $f(x) \rightarrow \infty$ as $x \rightarrow \infty$, such that

$$\mathcal{M}(\omega_\varphi) \geq f(I(\varphi)),$$

for all metrics $\omega_\varphi = \omega + \sqrt{-1}\partial\bar{\partial}\varphi$.

In [11, 12] it was used that in the toric case uniform K-stability is known to imply the properness of the Mabuchi energy by the work of Donaldson [7] and Zhou–Zhu [25], and moreover this is a condition that can be checked in certain cases. There are also other, non-toric, examples where properness of the Mabuchi energy is known, to which our result can be applied. For instance when M admits a positive Kähler–Einstein metric and has no holomorphic vector fields then the Mabuchi energy is proper in $c_1(M)$ (see Tian [22] and Phong–Song–Sturm–Weinkove [15]). If $c_1(M) = 0$, then the same is true in every Kähler class. If $c_1(M) < 0$, then the work of Song–Weinkove [19] gives an explicit neighborhood of the class $-c_1(M)$, where the Mabuchi energy is proper (see also [3], [24], [8]).

2. Proofs of the results. We can now proceed to the proof of Theorem 1.

PROOF OF THEOREM 1. We argue by contradiction. Given a constant $K > 0$, suppose that there does not exist a suitable $C > 0$ as in the statement of the proposition. We can then choose a sequence of smooth functions $\varphi_k \in PSH(M, \omega)$, such that $\omega_k = \omega + \sqrt{-1}\partial\bar{\partial}\varphi_k$ satisfy

$$|Rm(\omega_k)| < K, \quad I(\varphi_k) < K,$$

but there is no C for which $\omega_k > C^{-1}\omega$ and $\|\omega_k\|_{C^{1,\alpha}(\omega)} < C$ for all k . We can assume that

$$\|\omega_k\|_{C^{1,\alpha}(\omega)} + \sup \Lambda_{\omega_k} \omega > k.$$

Without loss of generality we can modify each φ_k by a constant in order to let $\sup_M \varphi_k = 0$. A standard argument using Green's formula and the inequality $\Delta \varphi_k > -n$ yields

$$(2) \quad \int_M \varphi_k \omega^n > -C_1$$

for some constant C_1 . It then follows from the bound $I(\varphi_k) < K$, that

$$(3) \quad E(\varphi_k) = I(\varphi_k) - \int_M \varphi_k \omega^n < K + C_1.$$

Since each ω_k is in the fixed class $[\omega]$, we can choose a subsequence (also denoted by ω_k for simplicity) such that the ω_k converge to a limiting current $\omega_\infty = \omega + \sqrt{-1}\partial\bar{\partial}\varphi_\infty$ weakly. It then follows that $\varphi_k \rightarrow \varphi_\infty$ in L^1 , so (3) together with Proposition 5 imply that $\varphi_\infty \in \mathcal{E}^1(M, \omega)$. Now Proposition 4 implies that φ_∞ has vanishing Lelong numbers.

Let $x \in M$ and $\delta > 0$. Since $\nu(\varphi_\infty, x) = 0$, there exists an $r > 0$ such that

$$c_n r^{2-2n} \int_{B_r(x)} \sqrt{-1}\partial\bar{\partial}\varphi_\infty \wedge \omega^{n-1} < \delta.$$

By choosing r smaller we can assume that

$$c_n r^{2-2n} \int_{B_r(x)} \omega_\infty \wedge \omega^{n-1} < \delta.$$

By the weak convergence of ω_k to ω_∞ we can choose $N > 0$ such that

$$c_n r^{2-2n} \int_{B_r(x)} \omega_k \wedge \omega^{n-1} < \delta, \quad \text{for } k > N.$$

Then by choosing δ sufficiently small, we can ensure that for $k > N$ we have

$$r^{2-2n} \int_{B_r(x)} e(\varphi_k) \omega^n < \epsilon.$$

Proposition 3 then implies that

$$\sup_{B_{r/2}(x)} e(\varphi_k) < 4r^{-2}$$

for $k > N$. For each x we obtain a different radius r , but the balls $B_{r/2}(x)$ cover M , and so we can choose finitely many of them which still give an open cover. It follows that we can choose a large N , and small $r > 0$ such that

$$\Delta \varphi_k + n = e(\varphi_k) < 4r^{-2}$$

on all of M , for $k > N$. In particular there is a constant C_2 such that

$$\Delta\varphi_k < C_2 \quad \text{for all } k,$$

and this gives upper bounds

$$(4) \quad \omega_k < (1 + C_2)\omega$$

on the metrics ω_k . Using Green's formula again, together with the bound (2), we get a uniform C^0 bound on the φ_k .

To obtain further bounds on the metrics, we could proceed as in Chen–He [4] Theorem 5.1. More directly, let

$$(5) \quad F_k = \log \frac{\omega_k^n}{\omega^n},$$

and note that

$$\sqrt{-1}\partial\bar{\partial}F_k = Ric(\omega) - Ric(\omega_k).$$

The upper bound (4) on the metrics ω_k and the uniform curvature bound imply that

$$(6) \quad |\Delta F_k| < C_3$$

for some C_3 . Since

$$\int_M \frac{\omega_k^n}{\omega^n} \omega^n = \int_M \omega_k^n = \int_M \omega^n,$$

we must have $F_k(x) = 0$ for some $x \in M$. Using Green's formula and $\Delta F_k > -C_3$ as we did for (2) we obtain

$$\int_M F_k > -C_4$$

for some C_4 . Using this and $\Delta F_k < C_3$ in Green's formula we get $F_k > -C_5$. Finally this bound together with the upper bound (4) implies a uniform lower bound on the metric ω_k . This, together with the curvature bound implies $C^{1,\alpha}$ -bounds on the metric ω_k .

In sum we have obtained a uniform constant C such that $\omega_k > C^{-1}\omega$ and $\|\omega_k\|_{C^{1,\alpha}(\omega)} < C$ for all k . This contradicts our assumption, and proves the theorem. \square

EXAMPLE 7. Note that in the 1-dimensional case $I(\varphi)$ is simply the L^2 -norm of the gradient of φ :

$$I(\varphi) = \int_M \varphi(-\sqrt{-1}\partial\bar{\partial}\varphi) = \int_M \sqrt{-1}\partial\varphi \wedge \bar{\partial}\varphi = \int_M |\partial\varphi|_\omega^2 \omega.$$

We will show that in Theorem 1 one cannot replace this with an L^p norm of the gradient for $p < 2$.

Let $M = \mathbf{P}^1$, and ω be the Fubini–Study metric given in a coordinate chart by

$$\omega = \sqrt{-1}\partial\bar{\partial} \log(1 + |z|^2).$$

Let

$$\omega_\lambda = \sqrt{-1}\partial\bar{\partial} \log(1 + |\lambda z|^2) = \omega + \sqrt{-1}\partial\bar{\partial} \log \frac{\lambda^{-2} + |z|^2}{1 + |z|^2}.$$

This is also the Fubini–Study metric, just in different coordinates, so $|Rm(\omega_\lambda)| < C$ for a constant C independent of λ . On the other hand the metrics are not uniformly equivalent, since $\omega_\lambda(0) = \lambda^2\omega(0)$. We will see that nevertheless the gradients of the Kähler potentials are uniformly bounded in L^p for any $p < 2$.

The Kähler potentials are

$$\varphi_\lambda = \log(\lambda^{-2} + |z|^2) - \log(1 + |z|^2),$$

and so we can compute

$$|\partial\varphi_\lambda|_\omega^2 = \frac{|z|^2(1 - \lambda^{-2})^2}{(\lambda^{-2} + |z|^2)^2}.$$

In polar coordinates the integral of $|\partial\varphi_\lambda|^p$ with respect to ω is

$$2\pi \int_0^\infty \frac{r^{p+1}(1 - \lambda^{-2})^p}{(\lambda^{-2} + r^2)^p(1 + r^2)^2} dr.$$

If $\lambda \geq 1$, then

$$\frac{r^{p+1}(1 - \lambda^{-2})^p}{(\lambda^{-2} + r^2)^p(1 + r^2)^2} \leq \frac{r^{1-p}}{(1 + r^2)^2}.$$

If $p < 2$, then the right hand side is integrable, so we have a uniform bound on the L^p -norm of $|\partial\varphi_\lambda|$.

PROOF OF THEOREM 2. Given Theorem 1, the proof of Theorem 2 is along standard lines, as in Chen–He [4] for instance. We outline the main points. Crucially, the Mabuchi energy is decreasing along a solution ω_t of the Calabi flow:

$$\frac{d}{dt} \mathcal{M}(\omega_t) = - \int_M (S(\omega_t) - \hat{S})^2 \omega_t^n \leq 0.$$

The properness assumption on the Mabuchi energy then gives a uniform bound on $I(\varphi_t)$ along the flow. This together with Theorem 1 implies that the metrics along the flow are uniformly equivalent. At this point one can show that the Calabi energy

$$\int_M (S(\omega_t) - \hat{S})^2 \omega_t^2$$

decays exponentially fast to zero. The smoothing property of the flow implies uniform bounds on the derivatives of $S(\omega_t)$, so from the decay of the Calabi

energy we find that in fact $S(\omega_t) - \hat{S} \rightarrow 0$ in any C^k norm, exponentially fast. This proves the exponential convergence of the flow

$$\frac{\partial}{\partial t} \varphi_t = S(\omega_t) - \hat{S}.$$

The limit is necessarily a constant scalar curvature metric in the class $[\omega]$. \square

References

1. Aubin T., *Réduction du cas positif de l'équation de Monge–Ampère sur les variétés kählériennes compactes à la démonstration d'une inégalité*, J. Funct. Anal., **57**, No. **2** (1984), 143–153.
2. Calabi E., *Extremal Kähler metrics*, Seminar on Differential Geometry, pp. 259–290, *Ann. of Math. Stud.*, **102**, Princeton Univ. Press, Princeton N.J., 1982.
3. Chen X. X., *On the lower bound of the Mabuchi energy and its application*, Internat. Math. Res. Notices, No. **12** (2000), 607–623
4. Chen X. X., He W. Y., *On the Calabi flow*, Amer. J. Math., **130**, No. **2** (2008), 539–570.
5. Chen X. X., He W. Y., *The Calabi flow on toric Fano surfaces*, Math. Res. Lett., **17**, No. **2** (2010), 231–241.
6. Chruściel P. T. *Semi-global existence and convergence of solutions of the Robinson–Trautman (2-dimensional Calabi) equation*, Comm. Math. Phys., **137**, No. **2** (1991), 289–313.
7. Donaldson S. K. *Scalar curvature and stability of toric varieties*, J. Differential Geom., **62**, No. **2** (2002), 289–349.
8. Fang H., Lai M., Song J., Weinkove B., *The J-flow on Kähler surfaces: a boundary case*, [arXiv:1204.4068](https://arxiv.org/abs/1204.4068).
9. Guan D., *Extremal solitons and exponential C^∞ convergence of the modified Calabi flow on certain \mathbf{CP}^1 bundles*, Pacific J. Math., **233**, No. **1** (2007), 91–124.
10. Guedj V., Zeriahi A., *The weighted Monge–Ampère energy of quasisubharmonic functions*, J. Funct. Anal., **250**, No. **2** (2007), 442–482.
11. Huang H., *Convergence of the Calabi flow on toric varieties and related Kähler manifolds*, [arXiv:1207.5969](https://arxiv.org/abs/1207.5969).
12. Huang H., *Toric surfaces, K-stability and Calabi flow*, [arXiv:1207.5964](https://arxiv.org/abs/1207.5964).
13. Lelong P., *Fonctions plurisousharmoniques et formes différentielles positives*, Gordon & Breach, Paris–London–New York, 1968.
14. Mabuchi T., *K-energy maps integrating Futaki invariants*, Tohoku Math. J. (2), **38**, No. **4** (1986), 575–593.
15. Phong D. H., Song J., Sturm J., Weinkove B., *The Moser–Trudinger inequality on Kähler–Einstein manifolds*, Amer. J. Math., **130**, No. **4** (2008), 1067–1085.
16. Phong D. H., Sturm J., *On stability and the convergence of the Kähler–Ricci flow*, J. Differential Geom., **72**, No. **1** (2006), 149–168.
17. Ruan W. D., *On the convergence and collapsing of Kähler metrics*, J. Differential Geom., **52**, No. **1** (1999), 1–40.
18. Schoen R., Uhlenbeck K., *A regularity theory for harmonic maps*, J. Differential Geom., **17**, No. **2** (1982), 307–335.
19. Song J., Weinkove B., *On the convergence and singularities of the J-flow with applications to the Mabuchi energy*, Comm. Pure Appl. Math., **61**, No. **2** (2008), 210–229.

20. Székelyhidi G., *The Calabi functional on a ruled surface*, Ann. Sci. Éc. Norm. Supér. (4), **42**, No. **5** (2009), 837–856.
21. Székelyhidi G., *The Kähler–Ricci flow and K-polystability*, Amer. J. Math., **132**, No. **4** (2010), 1077–1090.
22. Tian G., *Kähler–Einstein metrics with positive scalar curvature*, Invent. Math., **130**, No. **1** (1997), 1–37.
23. Tosatti V., *Kähler–Ricci flow on stable Fano manifolds*, J. Reine Angew. Math., **640** (2010), 67–84.
24. Weinkove B., *On the J-flow in higher dimensions and the lower boundedness of the Mabuchi energy*, J. Differential Geom., **73**, No. **2** (2006), 351–358.
25. Zhou B., Zhu X., *Relative K-stability and modified K-energy on toric manifolds*, Adv. Math., **219**, No. **4** (2008), 1327–1362.

Received August 29, 2012

Department of Mathematics
University of Notre Dame
Notre Dame, IN 46615, USA
e-mail: gszekely@nd.edu